Integrated FIT and FC Lower GI Pathway **REVISED COVID PATHWAY Lower GI Symptoms LOW RISK HIGH RISK Colorectal Cancer Colorectal Cancer** Does not fulfil NICE Guidance (NG12) Fulfils NICE Guidance (NG12) see local pathway here Age ≥50 Age <50 Age ≤50 with fresh rectal Request FIT test at time of 2ww referral bleeding, no other symptoms (except patients who have rectal mass/rectal ulceration/lower abdominal mass) DO NOT DELAY REFERRAL Colorectal cancer not suspected On ICE select: GP Profiles / GP Tests / FC care pathway if diarrhoea predominant FIT/FC Integrated pathway / Yes patient symptoms **Routine Referral to** meets 2ww criteria (responsibility of **Colorectal Surgery** secondary care); Perform blood investigations (FBC, ferritin, U+E) FC ≥ 100 FC <100 FIT pathway see local pathway here 2ww Referral Repeat FC within 2-4 weeks FIT ≥10 FIT <10 FC 100-250 Primary care FC < 100 FC >250 management Clinical review after 2WW and review symptomatic treatment **Urgent Gastro Routine Gastro** Referral Referral ≥60y & symptomatic <60y & symptomatic **Routine Gastro**

Referral